

Drug Information Sheet("Kusuri-no-Shiori")

Internal
Revised: 07/2007

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Flavitan Syrup 0.3%

Active ingredient:Flavin adenine dinucleotide sodium

Dosage form:clear yellow syrup

Print on wrapping:



Effects of this medicine

This medicine is a Vitamin B2 preparation that works to maintain the metabolism of the body at a normal level. It is usually used for the prevention and treatment of various diseases that are caused by Vitamin B2 deficiency or metabolic disorder including stomatitis, eczema, etc.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
- In general, for adults, take 5 to 45 mg of flavin adenine dinucleotide per day in 1 to 3 divided doses. The dosage may be adjusted according to the patient's age and symptoms. One milliliter of this medicine contains 3 mg of flavin adenine dinucleotide. Strictly follow the instructions.
- If you miss a dose, skip the missed dose and continue your regular dosing schedule from the next dose onwards. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- Your urine may turn more yellow than usual. This is because this medicine is excreted into urine after being used in the body.
- Special caution should be taken because this medicine may stain your clothes when in contact. If it occurs, wash with soap and water.

Possible adverse reactions to this medicine

No pertinent entries.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of infants and young children. Store away from direct sunlight, heat and humidity.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.