

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 06/2011

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Starsis Tablets 90mg

Active ingredient: Nateglinide

Dosage form: pink tablet (φ: 8.1mm, thickness: 4.4mm)

Print on wrapping: (Face) 糖尿病の薬, 食直前に服用, スターシス 90, (Back) 糖尿病の薬, スターシス 90mg, house-mark



Effects of this medicine

This medicine suppresses postprandial hyperglycemia (elevation in the blood glucose level after a meal) by stimulating beta cells in the pancreas to promote insulin secretion.

It is usually used for the treatment of diabetes mellitus.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you ever experienced any allergic reaction (itch, rash etc.) to any medicine.
If you have ketosis, diabetic coma or precoma, insulin-dependent diabetes mellitus, renal disorder, or infections.
If you are in a pre- or post-operative state.
If you have any injury.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
- General dosage regimen: For adults, take 1 tablet (90mg of the active ingredient) at a time, 3 times daily, before each meal (within 10 minutes). If the effect is insufficient, the dosage may be increased up to 120mg at a time while observing your clinical course. As this preparation contains 90mg of the active ingredient per tablet, take this preparation when you take at least 90mg at a time according to the above dosage regimen. Strictly follow the instructions of your doctor/pharmacist.
- If you realize that you have missed a dose after starting a meal, skip the missed dose and take the usual dose before the next meal. DO NOT take double doses to make up for the missed dose. (It may cause hypoglycemic conditions).
- If you took too much of the medicine (more than ordered), check with your doctor/pharmacist. If you have dizziness or light-headedness due to lowering of blood glucose (hypoglycemia), take something that contains glucose, such as sugar or juice, or glucose if you are concomitantly taking an alpha-glucosidase inhibitor (acarbose or voglibose), before consulting with your doctor.
- Do not stop taking the medicine without the instructions of your doctor.

Precautions while taking this medicine

- Share better understanding with your family about your diabetes mellitus.
- Take meals regularly and avoid excessive eating and drinking.
- The medicine may cause dizziness or light-headedness due to hypoglycemia. Pay close attention when you drive a car, work at heights, or operate dangerous machinery.
- Take something that contains glucose, such as sugar or juice, if you have dizziness or light-headedness due to hypoglycemia. Take glucose, if you are concomitantly taking an alpha-glucosidase inhibitor (acarbose or voglibose).

Possible adverse reactions to this medicine

Common side effects are reported as below. If any of them occurs, check with your doctor/pharmacist: queasy, increased flatus, bloating, diarrhoea, abdominal pain, rash, itching, urticaria, erythema multiforme, headache, dizziness, malaise, etc.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- cold sweat, chills, palpitations [hypoglycemia]

- general malaise, loss of appetite, yellowing of the skin or whites of the eyes [liver function disorder, jaundice]
- chest pain with feeling of pressure or tightness, anginal pain, cold sweat [myocardial infarction]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of infants and children. Store away from direct sunlight, heat and humidity.
- Keep the drug in a press-through-package (PTP), and remove the drug from the PTP sheet right before use.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.