

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 04/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Harnal D Tablets 0.1mg

Active ingredient:Tamsulosin hydrochloride

Dosage form:white orally-disintegrating tablet, diameter: 7.5 mm, thickness: 3.3 mm

Print on wrapping:(Face) ハルナール D 0.1mg, 0.1mg, 口腔内崩壊錠, アステラス製薬, (Back) HarnalD 0.1mg, ハルナール D, 0.1mg, house-mark, house-mark+HA0.1, アステラス製薬



Effects of this medicine

This medicine reduces urethral pressure by its blocking activity on alpha 1 receptors in the urethra and prostate, thus improving impaired urination associated with prostatic hyperplasia.

It is usually used for the treatment of impaired urination associated with prostatic hyperplasia.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 2 tablets (0.2 mg of the active ingredient) at a time, once a day, after a meal. The dosage may be adjusted according to the age or symptoms. Strictly follow the instructions.
- Take the tablet with water or lukewarm water in the same way as you usually take tablets. Since the tablet promptly dissolves in saliva, you may place it on your tongue, moisten with saliva, crush it with your tongue and swallow it together with saliva.
- Do not crunch the tablet when you take it.
- Do not take the tablet without water when you are lying down.
- If you miss a dose, skip the missed dose and follow your regular dosing schedule from the next day. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- This medicine may cause dizziness. Pay close attention when you perform dangerous operations such as working at heights or driving a car.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include dizziness, swaying feeling, tachycardia, rash, gastric distress, nausea, vomiting, gastric pain, loss of appetite and general malaise. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- fainting, loss of consciousness [syncope/unconsciousness]
- general malaise, loss of appetite, yellowing of the skin and the white of eyes [liver dysfunction, jaundice]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.