

# Drug Information Sheet("Kusuri-no-Shiori")

Injection

Published: 05/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**Shinbit Injection 50mg

**Active ingredient:**Nifekalant hydrochloride

**Dosage form:**Injection

**Print on wrapping:**



## Effects of this medicine

This medicine works to relieve abnormal excitation of the heart by blocking the potassium channel of cardiac muscle, and consequently normalize the pulse.

It is usually used for the treatment of ventricular tachycardia and ventricular fibrillation

## Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.  
If you have long QT syndrome.
- If you are pregnant, possibly pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, inject this medicine intravenously. Electrocardiography are performed on a continuous basis during treatment with this medicine.  
This medicine may be injected intravenously and continuously to maintain the effect.
- Consult with your doctor as to the specific treatment period.

## Precautions while taking this medicine

## Possible adverse reactions to this medicine

The most commonly reported adverse reactions include proarrhythmia. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, see your doctor immediately.**

- dizziness, chest pain, palpitation [proarrhythmia]

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

## Storage conditions and other information

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.