

Drug Information Sheet("Kusuri-no-Shiori")

Injection

Published: 05/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Cibenol Intravenous Injection 70mg

Active ingredient: Cibenzoline succinate

Dosage form: Injection

Print on wrapping:



Effects of this medicine

This medicine works to normalize the pulse by blocking the sodium channel (membrane stabilizing action) of cardiac muscle as well as prolonging the action potential duration.

It is usually used for the treatment of tachyarrhythmia.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
If you have atrioventricular block, sinoatrial block, congestive heart failure, angle-closure glaucoma or difficulty in urination.
- If you are on dialysis.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, this medicine is injected intravenously over 2 to 5 minutes. Blood pressure measurement and electrocardiography are performed on a continuous basis during treatment with this medicine.
- Consult with your doctor as to the specific treatment period.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include hot flush, bradycardia, decreased blood pressure, palpitation, dizziness, nausea, thirst and numbness. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, see your doctor immediately.

- dizziness, chest pain, palpitation [arrhythmogenic activity]
- facial pallor, cold sweat, dizziness on standing up [shock, anaphylaxis]
- shortness of breath, general edema, cough [cardiac failure]
- general malaise, loss of appetite, yellowing of the skin and the whites of eyes [liver disorder due to circulatory failure, hepatic function disorder, jaundice]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.